

VACCINATION

WHEN TO GET AN ANIMAL VACCINATED



Puppies (0-1 year old)

CORE VACCINE
(**PARVOVIRUS, DISTEMPER,**
ADENOVIRUS, PARAINFLUENZA)

Vaccinate at 2 months, 3 months and 1 year of age

RABIES VACCINE
Vaccinate at 3 months and 1 year of age



Adults

CORE VACCINE
(**PARVOVIRUS, DISTEMPER, ADENOVIRUS,**
PARAINFLUENZA)

Vaccinate every 3 years
(if never vaccinated, vaccinate once, then 1 year after the first vaccine, then every 3 years)

RABIES VACCINE
Vaccinate every 3 years
(if never vaccinated, vaccinate once, then 1 year after the first vaccine, then every 3 years)

DISTEMPER

- Transmission: air
- Signs: coughing, eye problems, vomiting, diarrhea, weakness, lost of balance
- Prevention: vaccination, isolation of the sick animal

PARVOVIRUS

- Transmission: stools (will persist in the environment up to 6 months)
- Signs: weakness, loss of appetite, vomiting, diarrhea
- Prevention: vaccination, isolation of the sick animal, removal of the animal's stools and, if indoors, the surfaces should be cleaned with diluted bleach

RABIES

- Transmission: bite from an infected animal
- Signs: loss of appetite, salivation, change in behaviour, lost of balance
- Prevention: vaccination

HOW TO GET YOUR ANIMAL VACCINATED

Contact the vaccinator in your community to have your animal vaccinated. You can also contact the municipal office for more information.



Contact the Veterinary Help Line to seek free veterinary advice at 514 345-8521
ext. 0065 or nunavik-giv@medvet.umontreal.ca